

Bibliography --Osteoporosis

Avenell A, Gillespie WJ, Gillespie LD, O'Connell D. *Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis*. Cochrane Database of Systemic Reviews 2009, Issue 2, Art. No.: CD000227. DOI: 10.1002/14651858.CD000227.pub3

Bennell KL, Matthews B, Greig A, Briggs A, Kelly A, Sherburn M, Larsen J, Wark J, BMC Musculoskelet Disord 2010; 11(36); epub-. *Effects of an exercise and manual therapy program on physical impairments, function and quality of life in people with osteoporotic vertebral fracture: a randomized, single-blind controlled pilot trial*

Black D, et al. *Once-a-year infusion reduces osteoporosis fracture risk*. New England Journal of Medicine 2007; 356: 1809-22

Bonaiuto D, Shea B, Iovine R, Negrini S, Welch V, Kemper HHC, Wells GA, Tugwell P, Cranney A. *Exercise for preventing and treating osteoporosis in postmenopausal women*. Cochrane Database of Systematic Reviews 2002, Issue 3. Art. No.: CD000333. DOI: 10.1002/14651858.CD000333

Chien MY, Yang RS, Tsauo JY. *Home-based trunk strengthening exercise for osteoporotic and osteopenic postmenopausal women with fracture—a pilot study*. Clin Rehabil, 2005; 19(1): 28-36

Devereux K, Robertson D, Briffa NK. *Effects of a water-based program on women 65 years and over: a randomised controlled trial*. Aust J Physiother, 2005; 51 (2): 102-108

Haugenauer D, Shea B, Tugwell P, Wells GA, Welch V. *Fluoride for treating postmenopausal osteoporosis*. Cochrane Database of Systemic Reviews 2000, Issue 4. Art. No.: CD002825. DOI: 10.1002/14651858.CD002825

Homik J, Cranney A, Shea B, Tugwell P, Wells GA, Adachi J, Suarez-Almazor ME. *Bisphosphonates for steroid-induced osteoporosis*. Cochrane Database of Systemic Reviews 1999, Issue 1. Art. No.: CD001347. DOI:10.1002/14651858. CD001347

Homik J, Suarez-Almazor ME, Shea B, Cranney A, Wells GA, Tugwell P. *Calcium and vitamin D for corticosteroid-induced osteoporosis*. Cochrane Database of Systemic Reviews 1998, Issue 2. Art. No.: CD000952. DOI:10.1002/14651858.CD000952

Kamide N, Shiba Y, Shibata H. *Effects on balance, falls, and bone mineral density of a home-based exercise program without home visits in community-dwelling elderly women: a randomized controlled trial*. J Physiol Anthropol, 2009; 28(3): 115-122

Kemmler W, Engelke K, von Stengel S, Weineck J, Lauber D, Kalender WA. *Long-term four-year exercise has a positive effect on menopausal risk factors: the Erlangen Fitness Osteoporosis Prevention Study*. J Strength Cond Res, 2007; 21 (1): 232-239

Maddalozzo G, et al. *Resistance training matches HRT in preventing spinal osteoporosis*. Bone, 2007; 40: 1244-51

Madureira MM, Takayama L, Gallinaro AL, Caparbo VF, Costa Ra, Pereira RM. *Balance training program is highly effective in improving functional status and reducing the risk of falls in elderly women with osteoporosis: a randomized controlled trial*. Osteoporos Int 2007; 18(4): 419-425

Pomerantz EM, *Osteoporosis and the female patient*, Orthopedic Physical Therapy Clinics in North America, 1996, 5, (1), 71-84.

Preisinger E, Alacamlioglu Y, Pils K, Bosina E, Metka M, Schneider B, Ernst E, Br J Sports Med 1996; 30(3); 209-212. *Exercise therapy for osteoporosis: results of a randomized controlled trial*

Shea B, Wells GA, Cranney A, Zytaruk N, Griffith L, Hamel C, Ortiz Z, Peterson J, Tugwell P, Welch V. *Calcium supplementation on bone loss in postmenopausal women*. Cochrane Database of Systemic Reviews 2004, Issue 1. Art. No.: CD004526. DOI: 10.1002/14651858.CD004526.pub3

Sinaki M, Mikkelsen BA. *Postmenopausal spinal osteoporosis: flexion versus extension exercises*. Arch Phys Med Rehabil, 1984; 65 (10): 593-596

Sinaki M, Brey RH, Hughes CA, Larson DR, Kaufman KR. *Significant reduction in risk of falls and back pain in osteoporotic-kyphotic women through a Spinal Proprioceptive Extension Dynamic (SPEED) program*. Mayo Clin Proc, 2005; 80(7): 849-855

Tucker K, et al. *Cola consumption may lower BMD in postmenopausal women*. American Journal of Clinical Nutrition 2006; 84: 936-42

Wells GA, Cranney A, Peterson J, Boucher M, Shea B, Welch V, Coyle D, Tugwell P. *Etidronate for the primary and secondary prevention of osteoporotic fractures in postmenopausal women*. Cochrane Database of Systemic Reviews 2008, Issue 1. Art. No.: CD003376. DOI: 10.1002/14651858.CD003376.pub3

Wells GA, Cranney A, Peterson J, Boucher M, Shea B, Welch V, Coyle D, Tugwell P. *Risedronate for the primary and secondary prevention of osteoporotic fractures in postmenopausal women*. Cochrane Database of Systemic Reviews 2008, Issue 1. Art. No.: CD004523. DOI: 10.1002/14651858.CD004523.pub3

Wells GA, Cranney A, Peterson J, Boucher M, Shea B, Welch V, Coyle D, Tugwell P. *Alendronate for the primary and secondary prevention of osteoporotic fractures in postmenopausal women*. Cochrane Database of Systemic Reviews 2008, Issue 1. Art. No.: CD001155. DOI: 10.1002/14651858.CD001155.pub2

U.S. Department of Health and Human Services. *Bone Health and Osteoporosis: A Report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services, Office of the Surgeon General, 2004.

Bibliography—Female Athlete

Braam L, Knapen M, Guesens P, Brouns F, Vermeer C. *Factors Affecting Bone Loss in Female Endurance Athletes*. Am J of Sports Med, 2003; Vol. 31, No. 6: 0363-5465/103/3131-0889\$02.00/0

Elliot D, Moe E, Goldberg L, DeFrancesco C, Durham M, Hix-Small H. *Definition and Outcome of a Curriculum to Prevent Disordered Eating and Body-shaping Drug Use*. J of School Health, 2006; Vol. 76, No. 2, 67-73

Gilchrist J, Mandelbaum B, Melancon H, Ryan G, Silvers H, Griffin L, Watanabe D, Dick R, Dvorak J. *A Randomized Controlled Trial to Prevent Noncontact Anterior Cruciate Ligament Injury in Female Collegiate Soccer Players*. Am J of Sports Med, 2008; 36;1476, DOI:10.1177/0363546508318188

Grindstaff T, Hammill R, Tuzson A, Hertel J. *Neuromuscular Control Training Programs and Noncontact Anterior Cruciate Ligament Injury Rates in Female Athletes: A Numbers-Needed-to-Treat Analysis*. Journal of Athletic Training, 2006;41 (4); 450-456

Hewett TE, Lindenfeld TN, Riccobene JV, Noyes FR. *The effect of neuromuscular training on the incidence of knee injury in female athletes. A prospective study.* Am J Sports Med, 1999; 27(6): 699-706

Hewett TE, Stroupe AL, Nance TA, Noyes FR. *Plyometric training in female athletes. Decreased impact forces and increased hamstring torques.* Am J Sports Med, 1996; 24 (6):765-772

Holm I, Fosdahl MA, Friis A, Risberg MA, Mykebust G, Steen H. *Effect of neuromuscular training on proprioception, balance, muscle strength, and lower limb function in female team handball players.* Clin J Sport Med, 2004; 14(2): 88-94

Hurd WJ, Chmielewski TL, Snyder-Mackler L. *Perturbation-enhanced neuromuscular training alters muscle activity in female athletes.* Knee Surg Sports Traumatol Arthrosc., 2006; 14(1): 60-69

Kovacs EJ, Birmingham TB, Forwell L, Litchfield RB. *Effect of training on postural control in figure skaters: a randomized controlled trial of neuromuscular versus basic off-ice training programs.* Clin J Sport Med, 2004; 14(4): 215-224

Mandelbaum B, Silvers H, Watanabe D, Knarr J, Thomas S, Griffin L, Kirkendall D, Garrett, Jr W. *Effectiveness of a Neuromuscular and Proprioceptive Training Program in Preventing the Incidence of Anterior Cruciate Ligament Injuries in Female Athletes.* Am J of Sports Med, 2005, Vol. 33, No. 7, DOI: 10.1177/0363546504272261

Mehlenbeck R, Ward K, Klesges R, Vukadinovich C. *International Journal of Sport Nutrition and Exercise Metabolism*, 2004, 14; 18-29

Myer GD, Ford KR, McLean SG, Hewett TE. *The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics.* Am J Sports Med, 2006; 34(3): 445-455

Myer G, Ford K, Hewett T. *Rationale and Clinical Techniques for Anterior Cruciate Ligament Injury Prevention among Female Athletes.* Journal of Athletic Training, 2004; 39 (4):352-364

Myer GD, Ford KR, McLean SG, Hewett TE. *The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics.* Am J Sports Med, 2006; 34(3): 445-455

Myer GD, Ford KR, Palumbo JP, Hewett TE. *Neuromuscular training improves performance and lower-extremity biomechanics in female athletes.* J Strength Cond Res, 2005; 19(1): 51-60

Myklebust G, Engebretsen L, Braekken IH, Skjølberg A, Olsen OE, Bahr R. *Prevention of anterior cruciate ligament injuries in female team handball players: a prospective intervention study over three seasons.* Clin J Sport Med, 2003; 13(2): 71-78

Pasanen K, Parkkari J, Pasanen M, Hiilloskorpi H, Mäkinen T, Jarvinen M, Kannus P. *Neuromuscular training and the risk of leg injuries in female floorball players: cluster randomised controlled study.* BMJ 2008; 337 (a295):-

Paterno MV, Myer GD, Ford KR, Hewett TE. *Neuromuscular training improves single-limb stability in young female athletes.* J Orthop Sports Phys Ther, 2004; 34(6): 305-316

Silvers H, Mandelbaum B. *Prevention of anterior cruciate ligament injury in the female athlete.* Br J Sports Med 2007; 41(Suppl 1): i52-i59. DOI: 10.1136/bjsm.2007.037200

Vescovi JD, Canavan PK, Hasson S. *Effects of a plyometric program on vertical landing force and jumping performance in college women.* Phys Ther Sport, 2008; 9(4): 185-192